

# Comments on "Sudden future singularities"

A. Kwang-Hua CHU \*

P.O. Box 30-15, Shanghai 200030, PR China

## Abstract

We make comments on the letter by Barrow [*Class. Quantum Grav.* **21** (2004) L79] since we cannot observe a singularity for  $p(t)$  as  $t \rightarrow t_s$ . Instead, there might be a singularity for  $\rho(t)$  for  $t \sim 0$  case.

PACS numbers: 98.80.Cq, 98.80.Bp, 98.80.Jk

Barrow just presented an interesting letter which shows that a singularity can occur at a finite future time in an expanding Friedmann universe even when  $\rho > 0$  and  $\rho + 3p > 0$  [1]. He has identified conditions that are sufficient to avoid the appearance of this type of sp-curvature singularity at finite future time. These results can either be used as a more fine-grained method to search for finitetime singularities and evaluate their likelihood of occurrence with realistic fluids or, if such singularities are regarded as causal pathologies, to impose realistic conditions which exclude them by fiat.

Consider the Friedmann universe with expansion scale factor  $a(t)$ , curvature parameter  $k$ , and Hubble expansion rate  $H = \dot{a}/a$ ; then ( $8\pi G = c = 1$ ) the Einstein equations reduce to

$$3H^2 = \rho - \frac{k}{a^2}, \quad \dot{\rho} + 3H(\rho + p) = 0, \quad (1)$$

and

$$\frac{\ddot{a}}{a} = -\frac{\rho + 3p}{6}. \quad (2)$$

He first looked informally at whether it is possible for any type of singularity to develop in which a physical scalar becomes infinite at a finite future comoving proper time  $t_s$ , when, say,  $a(t) \rightarrow a(t_s) \neq 0$  or  $\infty$ , and  $H(t) \rightarrow H_s < \infty$  and  $H_s > 0$ .

Guided by these heuristic arguments, he then constructed an explicit example by seeking, over the time interval  $0 < t < t_s$ , a solution for the scale factor  $a(t)$  of the form  $a(t) = A + Bt^q + C(t_s - t)^n$ , where  $A > 0, B > 0, q > 0, C$  and  $n > 0$  are free constants to be determined.

He can obtain (Eq. (14) in [1])

$$\frac{\ddot{a}}{a} \rightarrow -\infty,$$

whenever  $1 < n < 2$  and  $0 < q \leq 1$ ; the solution exists on the interval  $0 < t < t_s$ . Hence, as  $t \rightarrow t_s$  we have  $a \rightarrow a_s$ ;  $H_s$  and  $\rho_s > 0$  (for  $3q^2(a_s - 1)^2 t_s^{-2} > -k$ ) are finite but  $p_s \rightarrow \infty$ .

---

\*Present Address : Department of Physics, Xinjiang University, 14, Shengli Road, Urumqi 830046, PR China.

When  $2 < n < 3$ , he noted that  $\ddot{a}$  remains finite but  $d^3a/dt^3 \rightarrow \infty$  as  $t \rightarrow t_s$  as  $p_s$  remains finite but  $\dot{p}_s \rightarrow \infty$ . By contrast, there is an initial all-encompassing strong-curvature singularity, with  $\rho \rightarrow \infty$  and  $p \rightarrow \infty$ , as  $t \rightarrow 0$ . However, from Eqs. (14) and (4) of [1], he can see that  $\rho$  and  $\rho + 3p$  remain positive.

After considering two approximate forms of  $a(t)$  as  $t \rightarrow 0$  and  $t_s$ , he then claimed that this specific family of solutions shows that it is possible for an expanding universe to develop a 'big-rip' singularity at a finite future time even if the matter fields in the universe satisfy the strong-energy conditions  $\rho > 0$  and  $\rho + 3p > 0$ .

The present author would like to make comments about his derivations (especially about the equation (4) and its subsequent variations in [1]) and his results, say, equations (11) or (13) in [1]. Our second remark could be induced by the first remark!

The first remark has been partially addressed, as Barrow noticed that, there are two independent Friedmann equations for the three quantities  $a$ ,  $p$  and  $\rho$ . In the absence of a relation between any two of them there is an unconstrained degree of freedom. If we introduce an equation of state  $p(\rho)$  which bounds the pressure by some well-behaved function of the density for example,  $p < C\rho$  with  $C > 0$ -then the pressure singularity at a finite future time is eliminated if we require  $\rho > 0$  and  $\rho + 3p > 0$ . An upper bound on the ratio  $p/\rho$  corresponds to a bound on signal propagation of waves associated with small changes in pressure.

The second remark will be demonstrated by my illustration of figures once we plot  $\ddot{a}(t)/a(t)[\propto p(t)]$  and  $\rho(t)$  with respect to  $t$  (from the equation (13) of [1]). As Barrow mentioned,  $1 < n < 2$  and  $0 < q \leq 1$ , thus we choose  $n = 1.5$ ,  $q = 0.8$ , and  $t_s = 10$ . But, the trouble is that  $a_s$  is unknown. We then try three cases :  $a_s = 1.00, 1.05, 50.00$ . We can easily observe from Fig. 1 that there might be a singularity for  $\rho(t)$  but not for  $p(t)$  as  $t \sim 0$  once  $a_s \sim 1$ . We cannot observe a singularity for  $\rho(t)$  or  $p(t)$  once  $t \rightarrow t_s$  which Barrow claimed in [1] that there will be a singularity for  $p(t)$ ! These can be demonstrated in Figs. 1, 2 and 3 for either  $t_s = 10$  or  $t_s = 100$  once  $t \rightarrow t_s$ .

Let us consider the expression of  $\ddot{a}(t)/a(t)$ , the important term is  $t_s - t = K t_s$ . If  $K \ll 1$ , then there will be no singularity for  $p(t)$  as  $t \rightarrow t_s$  once  $t_s^2 \times K^{2-n} \sim O(1)$ !

To conclude in brief, our remarks as demonstrated in Figs. 1, 2 and 3 could not only be valid to the false statement which Barrow made in [1] but also be possible be valid to his subsequent paper (quite similar works) in [2].

## References

- [1] Barrow JD 2004 *Class. Quantum Grav.* **21** L79.
- [2] Barrow JD 2004 *Class. Quantum Grav.* **21** 5619.




Fig. 1  $p(t)[\propto \ddot{a}(t)/a(t)]$  and  $\rho(t)$  vs.  $t$ . Here,  $n = 1.5$ ,  $q = 0.8$ ,  $t_s = 10.0$ . There might be a singularity for  $\rho(t)$  but not for  $p(t)$  as  $t \sim 0$  (case of  $a_s \sim 1$ ; but not for  $t \rightarrow t_s$ ).




Fig. 2  $p(t)[\propto \ddot{a}(t)/a(t)]$  and  $\rho(t)$  vs.  $t$ . Here,  $n = 1.5$ ,  $q = 0.8$ ,  $t_s = 100$ . There is no singularity for  $p(t)$  or  $\rho(t)$  once  $t \rightarrow t_s (\equiv 100)$ .




Fig. 3  $p(t)[\propto \ddot{a}(t)/a(t)]$  and  $\rho(t)$  vs.  $t$ . Here,  $n = 1.5$ ,  $q = 0.8$ ,  $t_s = 100$ . There might be no singularity for  $p(t)$  or  $\rho(t)$  once  $t \rightarrow t_s (\equiv 100)$ .